



**Environmental  
Monitoring for  
Radioactivity  
around Devonport  
Royal Dockyard**

**Annual Report**

**2019**

---

**REVISION RECORD**

Issue	Date	Description		Name	Signature
1	August 2020	First Issue	Author	J Vickers	
			Checked by	M Clark	
			Approved by	R Foran	

---

**DOCUMENT RESPONSIBILITY**

The Director Assurance is responsible for alterations to this document.

---

## **Executive Summary**

This report presents the findings of Devonport Royal Dockyard Limited's (DRDL's) environmental radioactivity monitoring programme during 2019. DRDL is contracted by the Ministry of Defence (MOD) to refit, repair and maintain nuclear powered submarines at Devonport Royal Dockyard (DRD). During this work, quantities of liquid, solid and airborne radioactive waste are generated. DRDL is permitted under the Environmental Permitting (England and Wales) Regulations 2016 to dispose of the radioactive waste generated at DRD.

Under Condition 3.2.1(b) of its radioactive waste disposal Permit (Reference 1), DRDL is required to "define, document and carry out an environmental monitoring programme". DRDL's programme consists of sediment, biota and river water sampling and radiation monitoring surveys at numerous locations in the area around DRD (e.g. Tamar Estuary, Rivers Plym, Tavy and Lynher) and High Volume Air Sampling within DRD.

This report specifies the programmes for both marine and airborne radioactivity monitoring/sampling, presents and discusses the results of the monitoring/sampling and makes conclusions, comparing the results with independent monitoring programmes, regulatory dose limits, constraints and levels and with natural background radiation doses within the United Kingdom (UK).

Cobalt-60 (the radionuclide of most radiological significance in discharges from DRD) was not detected in any of the samples taken during 2019. The calculated annual effective radiation dose from DRDL's discharges of radioactive waste from DRD was less than 3  $\mu\text{Sv}$  per annum which equates to below 0.3% of the UK legal and International Commission for Radiological Protection (ICRP) dose limit of 1,000  $\mu\text{Sv}$  per annum for members of the public. The results of DRDL's environmental radioactivity monitoring programme are also comparable with the findings of the independent monitoring programmes reported in the Radioactivity in Food and the Environment (RIFE) reports, which state that the dose to members of the public from DRDL's radioactive discharges is less than 5  $\mu\text{Sv}$ . For comparison, the annual average background radiation dose in the UK is 2,700  $\mu\text{Sv}$ .

Based on the results in this report, and the continued downward trend in river discharges, it is concluded that DRD continues to be a site of low radiological significance.

## **CONTENTS**

### **EXECUTIVE SUMMARY**

- 1.0 INTRODUCTION
- 2.0 MONITORING PHILOSOPHY
- 3.0 THE MARINE ENVIRONMENTAL MONITORING PROGRAMME
- 4.0 MARINE ENVIRONMENTAL MONITORING RESULTS
- 5.0 DISCUSSION OF MARINE RESULTS
- 6.0 LIQUID RADIOACTIVE WASTE DISCHARGES
- 7.0 ENVIRONMENTAL MONITORING FOR AIRBORNE RADIOACTIVITY
- 8.0 CALCULATION OF RADIATION EXPOSURE
- 9.0 CONCLUSION
- 10.0 REFERENCES
- 11.0 GLOSSARY OF TERMS

### **ATTACHMENTS**

- 1 Marine environmental monitoring programme
- 2 Environmental monitoring programme for airborne radioactivity

### **TABLES**

- 1 Discharges of radioactive liquid waste to the Hamoaze during 2019
- 2 Gamma radiation dose rates 1m above inter-tidal sediment
- 3 Gamma radiation dose rates in contact with the strand line
- 4 Cobalt-60 in sediment samples
- 5 Cobalt-60 in seaweed samples
- 6 Cobalt-60, carbon-14 and tritium in shellfish samples
- 7 Cobalt-60, carbon-14 and tritium in fish samples
- 8 Cobalt-60 and tritium in river water samples
- 9 Gamma spectrometry and gross beta High Volume Air Sample results
- 10 Cobalt-60 and tritium rainwater results
- 11 A summary of the average concentrations of natural radioactivity found within the inter-tidal region in the UK
- 12 Average marine environmental monitoring results

### **GRAPHS**

- 1. Cobalt-60 and other radionuclides (excluding tritium) discharged per year into the Hamoaze via the existing dedicated pipeline.
- 2. Tritium activity discharged per year into the Hamoaze via the existing dedicated pipeline.

### **FIGURES**

- 1. Comparison of radiation doses.

## 1.0 INTRODUCTION

1.1 Devonport Royal Dockyard Limited (DRDL) is contracted by the Ministry of Defence (MOD) to refit, repair and maintain nuclear powered submarines at Devonport Royal Dockyard (DRD). During this work, quantities of liquid, solid and airborne radioactive waste are generated. DRDL is part of the Marine Sector of Babcock International Group plc. DRDL is Licensed by the Office for Nuclear Regulation (ONR) and Authorised by the Defence Nuclear Safety Regulator (DNSR) to operate DRD and is permitted by the Environment Agency under the Environmental Permitting (England and Wales) Regulations 2016 (EPR2016) to dispose of solid, liquid and gaseous radioactive waste generated at DRD.

1.2 DRDL's Permit (Reference 1) imposes limits and conditions relating to the methods by which liquid radioactive waste is disposed, the quantities of radioactivity permitted to be discharged, the samples to be taken and the records to be kept. Condition 3.2.1(b) of DRDL's Permit requires DRDL to "define, document and carry out an environmental radioactivity monitoring programme". Requirements for compliance with this permit condition are detailed in DRDL's management arrangements, which specifies that the results of DRDL's environmental radioactivity monitoring programme shall be reported quarterly and annually to the Environment Agency.

1.3 The purpose of DRDL's environmental radioactivity monitoring programme is to determine the radiation exposure of members of the public as a result of discharges of liquid radioactive waste from DRD. This report covers the results of monitoring/sampling carried out outside of DRD as well as High Volume Air Sampling (HVAS) within DRD and from the results assesses the potential exposure of the most exposed members of the public. It does not address radiation exposure of Company employees as this is covered by

other occupational monitoring programmes within DRD.

1.4 This annual report covers the environmental radioactivity monitoring programme and results for the period 1 January to 31 December 2019.

1.5 Defence Science and Technology Laboratory (Dstl) Chemical, Biological and Radiological Division (referred to in this Report as 'Dstl') publishes an annual report detailing its marine environmental radioactivity monitoring surveys at nuclear submarine berths (Reference 2). This report also includes the results of the quarterly monitoring surveys carried out by DRDL.

1.6 Independent radiological monitoring of food and the environment is also carried out around the dockyard through programmes sponsored by the Environment Agency and the Food Standards Agency. These monitoring programmes provide an independent check on the effects of discharges made by users of radioactive materials in the UK, and ensure that any radioactivity present in food and the environment does not compromise public health.

1.7 The results of monitoring around all major nuclear sites in the United Kingdom, including those from monitoring carried out around DRD, are published in the series of annual reports titled 'Radioactivity in Food and the Environment (RIFE)', the most recent of which is Reference 3. The RIFE Report contains results of radiological monitoring carried out by the UK environment agencies and Food Standards Agency.

1.8 Although some differences between the results can be expected because of the variability associated with biological systems and the uncertainties which are inherent in the measurement of radioactivity at the low levels present in the environment, the results presented in RIFE are generally consistent with those measurements made by DRDL, (see Section 4).

## **2.0 MONITORING PHILOSOPHY**

2.1 DRDL's radioactive waste stream contains primarily cobalt-60, tritium and carbon-14, with other radionuclides of lower radiological significance. Table 1 details the activity limits permitted for discharge to the Hamoaze and the activities discharged during 2019.

2.2 DRDL's monitoring philosophy is focused on the detection of cobalt-60, since in radiological terms cobalt-60 is recognised as being the most significant radionuclide in discharges from DRD. This is due to its physical characteristics, such as its relatively long half-life (i.e. 5.3 years), its emission of high energy gamma radiation and its tendency to adsorb onto marine sediments.

2.3 Further monitoring is carried out for tritium and carbon-14. Although the greatest quantity of activity discharged is tritium, this radionuclide, which also occurs naturally in the environment, is of less radiological significance than cobalt-60 primarily because it emits low energy beta radiation, is readily dispersed and diluted in the environment and, if an intake were to occur, is relatively quickly excreted from the body.

2.4 Carbon-14 is present in discharges from the dockyard and also occurs naturally in the environment. It has a long half-life and decays through emission of a low energy beta particle.

## **3.0 THE MARINE ENVIRONMENTAL MONITORING PROGRAMME**

3.1 In order to monitor the effect of liquid radioactive waste discharges on the marine environment, a radiation monitoring and sampling survey is undertaken each calendar quarter. The survey programme, which is detailed in Attachment 1, is carried out by DRDL in conjunction with Dstl. The monitoring programme has been agreed with the Environment Agency, MOD, Dstl and the local authorities.

3.2 Attachment 1 provides a description of DRDL's marine environmental monitoring programme. The following types of measurement are made at up to thirteen points centred around the Tamar estuary (measurements are also taken in the rivers Plym, Tavy and Lynher):

- Gamma radiation dose rates at 1 metre above the ground. (A measurement of the gamma radiation in air resulting from natural sources and from any contamination present in the sediment.)
- Gamma radiation dose rates in contact with the strand-line. (Measurements of contact dose rate along the high tide mark where there is a visible accumulation of items deposited on the shoreline.)
- High resolution gamma spectrometry analysis of samples of sediment, seaweed, shellfish, fish and river water. (Identifies and assesses concentrations of gamma emitting radionuclides present in the sample).
- Liquid scintillation analysis of shellfish and fish to identify the beta emitting radionuclides tritium and carbon-14. Samples are analysed for total tritium to ensure that any tritiated water present in each sample is included in the result, in addition to any organically bound tritium.

## **4.0 MARINE ENVIRONMENTAL MONITORING RESULTS**

4.1 The measurements made during 2019 are summarised in Tables 2 to 8. Where 'ND' is used in the tables, this means 'not detected' (i.e. the activity level within each such sample was below the limit of detection (LoD) for the analysis of that sample). The generic limit of detection used by DRDL/Dstl for cobalt-60 in samples is 1.0 becquerel per kilogram ( $Bqkg^{-1}$ ). Limits of detection for carbon-14 and tritium in biota samples are 50  $Bqkg^{-1}$  and 20  $Bqkg^{-1}$  respectively.

## **5.0 DISCUSSION OF MARINE RESULTS**

5.1 Whilst the DRDL marine environmental radioactivity monitoring programme focuses on the detection and measurement of cobalt-60, the gamma spectrometry analyses of the sediment, seaweed, shellfish, fish and river water routinely search for the presence of other significant gamma emitting radionuclides. The only gamma emitting radionuclides detected in 2019 are those which occur naturally and trace levels of the fission product caesium-137 mainly attributable to atmospheric weapons testing and the Chernobyl accident in 1986.

5.2 The average environmental gamma dose rate measured at a height of 1 metre around the inter-tidal area during 2019 was 0.086 microgray per hour ( $\mu\text{Gyh}^{-1}$ ), with individual results ranging from 0.062 to 0.109  $\mu\text{Gyh}^{-1}$ , (see Table 2 and Table 12). These gamma radiation dose rates give no environmental radiological concern and are comparable with published figures of radiation dose rates measured independently around Devonport (Reference 3), which are in the region of 0.074 to 0.10  $\mu\text{Gyh}^{-1}$ .

5.3 No cobalt-60 was detected in any of the marine samples collected by DRDL during 2019. A low level of carbon-14 was measured in fourteen biota samples and six samples had a low level of tritium.

5.4 The results are comparable with results for these radionuclides in biota detected at other UK sites (Reference 3) and are of low radiological significance.

5.5 Published data on natural radioactivity concentrations is summarised in Table 11 for comparison with the results obtained at Devonport. The average environmental monitoring results for Devonport in 2019 are presented in Table 12.

## **6.0 LIQUID RADIOACTIVE WASTE DISCHARGES**

6.1 All discharges to the Hamoaze during the reporting period were below DRDL's permitted limits, which are provided in Table 1. Graphs 1 and 2 show the levels of radioactivity discharged to the Hamoaze via the dedicated pipeline since 1987.

6.2 The volume of discharges to the river also varies depending on the work packages being undertaken. There was a decrease in the volume of discharges to the Hamoaze during the reporting period, from 397.8 m<sup>3</sup> in 2018 to 338.9 m<sup>3</sup> in 2019.

## **7.0 ENVIRONMENTAL MONITORING FOR AIRBORNE RADIOACTIVITY**

7.1 DRDL's airborne environmental monitoring programme includes High Volume Air Sampling (HVAS) and rainwater sampling and analysis (see Attachment 2). While DRDL have continued HVAS and rainwater sampling in 2019, it does not form part of the statutory monitoring programme, having been removed in January 2019 following discussion with the EA.

7.2 The results from the airborne monitoring programme carried out from January to December are summarised in Tables 9 and 10. Where ND is used in the Tables, this means 'not detected' (i.e. the activity levels were below the limit of detection).

7.3 No tritium or cobalt-60 was detected in any of the HVAS or rainwater samples. The positive gross beta and beryllium-7 results from the HVAS programme are attributed to naturally occurring radionuclides.

7.4 In accordance with the approach taken by the Environment Agency and Food Standards Agency (Reference 3) the levels of radionuclides detected in the airborne radioactivity samples are so low that radiological assessment of internal exposure from airborne radioactivity is not considered necessary.

## **8.0 CALCULATION OF RADIATION EXPOSURE**

8.1 In order to interpret the results of the marine environmental radioactivity monitoring programme in terms of radiation dose to members of the public, it is necessary to identify the most significant pathway, or combination of pathways, for radiation exposure and the group of people who are most likely to be exposed to the radiation from this (these) exposure pathway(s) (i.e. the 'representative individual'). A Radiological Habits Survey was carried out by Cefas in 2017 on behalf of the Environment Agency, Food Standards Agency and the ONR (Reference 5). Habits Surveys identify the occupancy levels, activities and eating patterns (including types of food, consumption rates, etc.) of the local population. This information is then used by the regulators to review the adequacy of the environmental monitoring programmes for the area.

8.2 Around Devonport, the representative individual for exposure from the marine pathways includes a member of the local community who resides or works on the mud flats and who eats locally caught seafood (Reference 3).

8.3 The quantity of main interest in terms of the radiological effect is the effective dose to the representative individual. The unit of effective dose is the sievert (Sv). Since environmental gamma dose rates measured at a height of one metre are generally indistinguishable from the natural background levels (Reference 4) it is not possible to use these values directly to calculate the radiation doses resulting from liquid effluent discharges.

8.4 Consequently, an alternative method is used to determine the external dose. Models defined in Reference 6 enable the environmental dose rate resulting from a particular concentration of radionuclide and type of sediment to be calculated. Using information provided in Reference 6, the effective dose rate above

sediment uniformly contaminated with cobalt-60 is calculated to be 0.6 nanosievert per hour ( $\text{nSvh}^{-1}$ ) per  $\text{Bqkg}^{-1}$ . As no cobalt-60 was detected in the sediment samples, it is conservatively assumed that the activity is present, but just at the limit of detection, and the LoD value (i.e.  $1 \text{ Bqkg}^{-1}$ ) is assigned. The maximum annual effective dose to the representative individual from external exposure over intertidal areas, for occupancy of 2100 hours per year over mud, is therefore calculated, using the models given in Reference 6, to be  $1.3 \mu\text{Svy}^{-1}$ .

8.5 The internal dose to members of the local community who consume locally caught fish and shellfish (the main contributors to the total internal dose) is assessed using an average of results obtained from the sampling programme. The dose contribution from a radionuclide in each food type was calculated using the average concentration for that radionuclide (the average is calculated by pessimistically assigning the LoD to samples where no activity was detected, along with any positive results). This average specific activity, together with data on the consumption rates and dose per unit intake from ingestion of seafood as provided in References 5, 7 and 8 was used to calculate the internal dose. The dose to the representative individual from internal exposure to cobalt-60, tritium and carbon-14 is calculated to be less than  $1 \mu\text{Svy}^{-1}$ .

8.6 The annual effective dose to the most exposed members of the local community from external and internal exposure as a result of radioactive liquid discharges from DRD is therefore assessed to be less than  $3 \mu\text{Svy}^{-1}$ .

## **9.0 CONCLUSION**

9.1 The results of DRDL's environmental radioactivity monitoring programme show that Devonport continues to be a site of low radiological significance.

9.2 The annual effective dose of less than  $3 \mu\text{Svy}^{-1}$  is:

- less than 0.3% of the current International Commission on Radiological Protection (ICRP) principal limit of exposure for members of the public of  $1 \text{ mSvy}^{-1}$  (Reference 9);
- less than 1% of the dose constraint of  $500 \mu\text{Svy}^{-1}$  for discharges from a single site (Reference 10);
- less than 2% of the dose constraint of  $300 \mu\text{Svy}^{-1}$  for discharges from a single source (Reference 10);
- well below the 'threshold of optimisation level' of  $20 \mu\text{Svy}^{-1}$  referred to in Command 2919 (Reference 11)

9.3 Exposure to natural radiation within the UK averages some  $2200 \mu\text{Sv}$  per person per annum with annual variations of several thousand  $\mu\text{Sv}$  in different parts of the UK, notably Devon and Cornwall (Reference 12). This can help to put the magnitude of the dose to the representative individual for Devonport (less than  $3 \mu\text{Svy}^{-1}$ ) into perspective (see Figure 1).

## 10.0 REFERENCES

- 1 DRDL Permit for the Receipt and Disposal of Radioactive Waste, Number EPR/BB3098DX
- 2 Dstl Report, "Marine Environmental Radioactivity Surveys at Nuclear Submarine Berths 2018".
- 3 "Radioactivity in Food and the Environment, 2018 (RIFE-24)", Food Standards Agency, Scottish Environment Protection Agency, Environment Agency and the Northern Ireland Environment Agency.
- 4 Aquatic Environment Monitoring Report No 45, "Radioactivity in Surface and Coastal Waters of the British Isles 1994" by W C Camplin, MAFF Directorate of Fisheries Research, (now Cefas), Lowestoft (1995).
- 5 Cefas Radiological Habits Survey: Devonport, 2017. K. J. Moore, F. J. Clyne, B. J. Greenhill and K. Clarke (2018).
- 6 Simple models for Prediction of External Radiation Exposure from Aquatic Pathways" by G J Hunt; Radiation Protection Dosimetry Vol. 8 No. 4 p 215-224 (1984).
- 7 "Generalised Habit Data for Radiological Assessments" by C A Robinson; NRPB-M636, 1996."
- 8 International Commission on Radiological Protection, Age-dependent Doses to Members of the Public from Intake of Radionuclides: Part 5, Compilation of Ingestion and Inhalation Dose Coefficients, ICRP Publication 119, 2012.
- 9 International Commission on Radiological Protection (ICRP); ICRP Publication 103; "2007 Recommendations of the International Commission on Radiological Protection". Annals of the ICRP Vol. 37 no. 2-4 (2007).
- 10 The Environmental Permitting (England and Wales) Regulations 2016, Schedule 23, Part 4.
- 11 "Review of Radioactive Waste Management Policy - Final Conclusions". Cmnd. 2919 (1995).
- 12 PHE-CRCE-026: Ionising Radiation Exposure of the UK Population: 2010 review. W Oatway et al. April 2016

---

## 11.0 GLOSSARY OF TERMS AND ABBREVIATIONS

Activity	Attribute of an amount of radionuclide. Describes the rate at which transformations occur in it – each transformation being associated with the emission of radiation. Unit becquerel, symbol Bq (see below). 1Bq = 1 transformation per second.
Beta Particle	An energetic electron emitted by some radionuclides.
Bq	The unit of radioactivity is the becquerel (Bq). In this report, cobalt-60 concentrations are expressed in terms of becquerel per kilogram of sample material (Bqkg <sup>-1</sup> ).
Cefas	Centre for Environment, Fisheries and Aquaculture Science.
DRD	Devonport Royal Dockyard. (Note: The overall site at Devonport consists of two parts; the Naval Base and Devonport Royal Dockyard.)
DRDL	Devonport Royal Dockyard Limited.
Dose rate	The rate at which radiation dose is received.
Dstl	Defence Science and Technology Laboratory (Chemical, Biological and Radiological Division).
Effective dose	An indicator of the effects of radiation on the body as a whole when different body tissues are exposed to different amounts and types of radiation (measured in Sievert (see below)).
Electron	A negatively charged sub-atomic particle.
EPR 2016	Environmental Permitting (England and Wales) Regulations 2016.
Gamma Radiation	Electromagnetic radiation, without mass or charge, emitted by some radionuclides.
G	giga (10 <sup>9</sup> )
Gray (Gy)	Quantity of energy imparted by ionising radiation to unit mass of matter such as tissue. This quantity is the absorbed dose and the standard international (SI) unit for absorbed dose is the Gray. 1 Gy = 1 joule per kilogram.
ICRP	International Commission on Radiological Protection.
Limit of Detection (LoD)	The limit at which it may be stated with a certain degree of statistical confidence (dependent upon the confidence limit, 95% or 99%) that activity has been detected.
MOD	Ministry of Defence.
M	mega (10 <sup>6</sup> )
m	milli (10 <sup>-3</sup> )

μ	micro ( $10^{-6}$ )
n	nano ( $10^{-9}$ )
ND	Not Detected.
PHE (RPD)	Public Health England (Radiation Protection Division). Formerly National Radiological Protection Board, then Health Protection Agency.
Representative Individual	A hypothetical individual receiving a dose that is representative of the most exposed individuals in the population (formerly the 'Critical Group').
Sievert (Sv)	The unit of Effective Dose (see above) obtained by multiplying the absorbed dose (see gray) by a factor to allow for the relative biological effectiveness of the various ionising radiations (for gamma, beta and X-rays the factor is 1).

---

## Attachment 1

### Marine environmental monitoring programme

#### 1.1 *Gamma Dose Rate*

Gamma dose rate measurements are made quarterly 1 metre above inter-tidal sediment at:

Royal Albert Bridge (east)	Royal Albert Bridge (west)	Bull Point
Wearde Quay	Thanks Park (Torpoint)	Slipway (Tamar St)
Stonehouse	Torpoint (south)	

and annually at:

Calstock	St Germans Quay	Mount Gould
Mount Jessop	Bovisand Bay	

Contact dose rate measurements are made bi-annually along strand lines at:

Bull Point	Wearde Quay
------------	-------------

#### 1.2 *Sediment*

Samples of inter-tidal surface sediment (scraped from the top few mm) are collected quarterly at:

Bull Point	Wearde Quay	Stonehouse	Torpoint (south)
------------	-------------	------------	------------------

and annually at:

Calstock	St Germans Quay	Mount Jessop	Bovisand Bay	Mount Gould
----------	-----------------	--------------	--------------	-------------

#### 1.3 *Seaweed*

Samples of *Fucus* seaweeds (usually '*Fucus vesiculosus*') are collected bi-annually at:

Bull Point	Wearde Quay	Stonehouse	Torpoint (south)
------------	-------------	------------	------------------

and annually at:

Mount Jessop	St Germans Quay	Mount Gould	Bovisand Bay
--------------	-----------------	-------------	--------------

The top 5 cm (i.e. recent growth) of samples, rinsed free of sediment, is analysed.

#### 1.4 *Fish and Shellfish*

Fish caught in Plymouth Sound (within the Breakwater) were obtained from local suppliers during the summer.

Samples of mussels '*Mytilus edulis*' of the largest size available are collected quarterly at Royal Albert Bridge (East) and Wearde Quay when available in sufficient numbers.

Mussels are prepared and the flesh extracted as if for eating. Only the mussel flesh is analysed.

### **1.5 River Water**

1 litre samples of river water are collected quarterly from the River Tamar at:

Royal Albert Bridge (East)      Torpoint (South)

All intertidal survey points are shown in Map 1.

**Map 1 – Marine environmental monitoring programme – intertidal survey locations**



**Attachment 2**

**Environmental monitoring programme for airborne radioactivity**

	<b>Description</b>	<b>Frequency of Analysis</b>	<b>Analysis</b>
<b>High Volume Air Sampler (HVAS)</b>	A HVAS is located within DRD. It operates at a flow rate of 1000 litres per minute. The filter papers are changed periodically and bulked together for analysis.	Quarterly	Gamma spectrometry and gas flow proportional counting.
<b>Rainwater Sampler</b>	Rainwater is collected at the dockyard. Once every quarter a one litre sample is taken from the rainwater collected and sent away for analysis.	Quarterly	Gamma spectrometry and liquid scintillation for tritium.



**The High Volume Air Sampler at the Dockyard**

**TABLE 1**

**Discharges of radioactive liquid waste to the Hamoaze during 2019**

<b>Radionuclide</b>	<b>Discharge Limit (GBq)</b>	<b>Discharges (GBq)</b>	<b>% of limit</b>
Cobalt 60	0.80	0.002	0.3
Tritium	700	4.98	0.7
Carbon-14	1.70	0.026	1.5
Other Radionuclides <sup>1</sup>	0.30	0.003	1.1

- Notes:
1. Total activity of all other radionuclides.
  2. The volume of effluent discharged to the Hamoaze during 2019 was 338.9 m<sup>3</sup>.

**Marine environmental monitoring results for 2019**

**Table 2**  
**Gamma radiation dose rates at 1 metre above inter-tidal sediment ( $\mu\text{Gy}\cdot\text{h}^{-1}$ )**

Location [Point Number]	2019 Doserates ( $\mu\text{Gy}/\text{h}$ )			
	January	April	July	October
Royal Albert Bridge (East) [1]	0.095	0.09	0.096	0.079
Thankes Park (Torpoint) [4]	0.082	0.088	0.097	0.092
Wearde Quay (Saltash) [5]	0.089	0.088	0.094	0.081
Slipway (Tamar Street) [6]	0.067	0.068	0.080	0.072
Royal Albert Bridge (West) [8]	0.088	0.09	0.084	0.087
Torpoint (South) [10]	0.09	0.098	0.096	0.103
Stonehouse [14]	0.068	0.066	0.073	0.062
Mount Gould [22]			0.097	
Mount Jessop [26]			0.102	
St Germans Quay [29]			0.092	
Bovisand Bay [30]			0.069	
Calstock [32]			0.109	
Bull Point [40]	0.088	0.086	0.085	0.083

Notes: (1) Measurements were made using a RadEye GX with geiger-müller probe on a 1m tripod stand.

(2) Points 22, 26, 29, 30 and 32 are only surveyed once a year.

**Table 3**  
**Gamma radiation dose rates in contact with the strand line ( $\mu\text{Gyh}^{-1}$ )**

Location	2019 Contact Dose rates ( $\mu\text{Gyh}^{-1}$ )			
	January		July	
	Maximum	Average	Maximum	Average
Wearde Quay (Saltash)	0.126	0.119	0.114	0.107
Bull Point	0.125	0.119	0.117	0.112

Notes: (1) Measurements were made using a RadEye GX with the geiger-müller probe placed in contact with the strandline.

**Table 4**  
**Cobalt-60 in sediment samples ( $\text{Bqkg}^{-1}$ ) (dry weight)**

Location	2019 Cobalt-60 Activity ( $\text{Bqkg}^{-1}$ )			
	January	April	July	October
Wearde Quay (Saltash)	ND	ND <sup>1</sup>	ND <sup>1</sup>	ND <sup>1</sup>
Torpoint (South)	ND	ND <sup>1</sup>	ND	ND <sup>1</sup>
Stonehouse	ND <sup>1</sup>	ND <sup>1</sup>	ND <sup>2</sup>	ND <sup>1</sup>
Mount Gould			ND <sup>2</sup>	
Mount Jessop			ND	
St Germans Quay			ND	
Bovisand Bay			ND <sup>2</sup>	
Calstock			ND	
Bull Point	ND <sup>1</sup>	ND <sup>1</sup>	ND	ND <sup>1</sup>

Note: All sediment samples were mud except for (1) mud/sand sample  
(2) sand sample

'ND' = Not Detected. Generalised limit of detection for all samples was  $1 \text{ Bqkg}^{-1}$ .  
Mount Gould to Calstock points are only surveyed once a year.

**Table 5**  
**Cobalt-60 in seaweed samples (Bqkg<sup>-1</sup>) (wet weight)**

Location	2019 Cobalt-60 Activity (Bqkg <sup>-1</sup> )	
	January	July
Wearde Quay (Saltash)	ND	ND
Torpoint (South)	ND	ND
Stonehouse	ND	ND
Mount Gould		ND
Mount Jessop		ND
St Germans Quay		ND
Bovisand Bay		ND <sup>1</sup>
Bull Point	ND	ND

Note: All seaweeds were fucus vesiculosus except for: (1) fucus serratus.  
'ND' = Not Detected. Generalised limit of detection for all samples was 1 Bqkg<sup>-1</sup>.

**Table 6 - Cobalt-60, Carbon-14 and Tritium in shellfish (M.edulis) samples (Bqkg<sup>-1</sup> Wet Weight) (2019)**

Location	Radionuclide	January	April	July	October	Limit of Detection <sup>2</sup>
Royal Albert Bridge (East)	Cobalt-60	ND	ND	ND	ND	1
	Carbon-14	28(9)	39(13)	17(7)	ND	50
	Tritium <sup>1</sup>	ND	ND	7(9)	17(13)	20
Wearde Quay	Cobalt-60	ND	ND	NA	NA	1
	Carbon-14	27(10)	48(15)	NA	NA	50
	Tritium <sup>1</sup>	ND	ND	NA	NA	20

- Notes: 1 Total Tritium.  
2 Actual limits of detection vary between samples. This can give rise to positive results which are less than the standard limits of detection required by the monitoring programme.  
3 Uncertainties of two standard deviations are shown in brackets.  
4 ND = Not Detected.  
5 NA = Not Available. Mussels were not available for collection

**Table 7**  
**Cobalt-60, Carbon-14 and Tritium in fish samples (2019)**  
**(Bqkg<sup>-1</sup> wet weight)**

<b>Fish Species</b>	<b>Cobalt-60</b>	<b>Carbon-14</b>	<b>Tritium</b>
Limit of Detection <sup>1</sup> (Bqkg <sup>-1</sup> )	1	50	20
Pollock	ND	24(10)	12(9)
Pollock	ND	21(10)	12(10)
Pollock	ND	19(9)	50(16)
Pollock	ND	23(9)	17(10)
Pollock	ND	19(9)	ND
Pollock	ND	29(11)	ND
Pollock	ND	39(13)	ND
Pollock	ND	24(10)	ND
Pollock	ND	24(10)	ND

Notes:

- 1 Actual limits of detection vary between samples. This can give rise to positive results which are less than the standard limits of detection required by the monitoring programme.
- 2 Uncertainties shown in brackets include all analytical uncertainties in accordance with ISO 17025 and provide a level of confidence of approximately 95% confidence level
- 3 ND = not detected

**Table 8  
Cobalt-60 and Tritium in river water samples (2019) (Bq l<sup>-1</sup>)**

Location	Radionuclide	January	April	July	October	Limit of Detection
Royal Albert Bridge (East)	Cobalt-60 Tritium	ND ND	ND ND	ND ND	ND ND	1 10
Torpoint (South)	Cobalt-60 Tritium	ND ND	ND ND	ND ND	ND ND	1 10

**Airborne environmental monitoring results for 2019**

**Table 9 Gamma spectrometry and Gross Beta high volume air sample results**

Measurement Period	Radioactivity Concentration (Bq)		
	Gross Beta*	Be-7	Co-60
Quarter 1	28(4)	124(20)	ND
Quarter 2	39(6)	43(9)	ND
Quarter 3	23(4)	187(24)	ND
Quarter 4	21(5)	329(52)	ND
Average Activity Concentration	173 $\mu\text{Bq m}^{-3}$	1.1 $\text{mBq m}^{-3}$	6 $\mu\text{Bq m}^{-3}$

**Notes:** Uncertainties shown in brackets include all analytical uncertainties in accordance with ISO 17025 and provide a level of confidence of approximately 95% confidence level  
 \_\_\_\_\_ 'ND' – Not Detected.

**Table 10 Cobalt-60 and Tritium rainwater results**

Measurement Period	Radioactivity Concentration (Bq l <sup>-1</sup> )	
	Tritium	Cobalt-60
Generic Limit of Detection	10	1
Quarter 1	ND	ND
Quarter 2	ND	ND
Quarter 3	ND	ND
Quarter 4	ND	ND

'ND' – not detected.

**Table 11**

**A Summary of the average concentrations of natural radioactivity found within the inter-tidal region in the UK.<sup>(1)</sup>**

Material	Total Beta Radioactivity Concentration (wet weight) <sup>(2)</sup>	
	Bqkg <sup>-1</sup>	Comments
Fish	40-100	Mostly K-40
Shellfish	40-100	Mostly K-40
Seaweed	200-600	Mostly K-40
Sand	200-400	K-40 and decay products of U and Th
Mud	700-1000	K-40 and decay products of U and Th

- Notes:
- (1) The information presented is taken from Reference 4.
  - (2) Except sediments for which dry weight concentrations apply.
  - (3) K-40 is naturally occurring potassium-40.

**Table 12**  
**Average marine environmental monitoring results for 2019**

Sample Type	No. of Samples	Mean Radioactivity Concentrations (Bq/kg) (Bq/l for water)				
		Co-60	Cs-137	K-40	H-3	C-14
Sediment	21	[1]	1.95	652	N/A	N/A
Seaweed	12	[1]	[1]	268	N/A	N/A
Mussels	6	[1]	[1]	44	16.33	28.67
Fish	9	[1]	[1]	93	21.11	25.00
River Water	8	[1]	[1]	[20]	[6.85]	N/A

<b>Mean Gamma Dose rate in air at 1m</b>	0.086 $\mu\text{Gyh}^{-1}$
<b>Mean Contact Gamma Dose rate along the strandline</b>	0.117 $\mu\text{Gyh}^{-1}$


Notes: (1) Values shown in square brackets indicate that there were no positive results in those samples. The figure given is the limit of detection. The respective LoD value for each radionuclide is assigned to all samples in which no activity is detected (ND), thus a pessimistic mean concentration is calculated.

(2) The samples of sediment, seaweed, river water, shellfish and fish contained some natural radioactivity predominately in the form of potassium-40 (K-40), which is present in all living, or once living, organisms and vegetation. Samples of sediment and seaweed also contained decay products of naturally occurring thorium and uranium.

(3) Very small quantities of caesium-137 (Cs-137) were detected in some environmental samples. This is attributed to atomic weapons test fall-out, discharges from other nuclear establishments and the Chernobyl accident.

(4) Results for sediment samples are dry weight concentrations, and results for biota are wet weight concentrations


**Graph 1 - Cobalt-60 and Other Activity (excluding Tritium) Discharged to the Hamoaze**



Notes:

- (1) The discharge of effluent for 1987 was recorded from June 1987 to December 1987 (i.e. after DRDL's first Radioactive Substances Act Authorisation was issued).
- (2) Prior to 1992 only gamma emitting radionuclides were included in 'Other Radionuclides'. From 1992 onwards this group includes both gamma and beta emitting radionuclides.
- (3) Carbon-14 was not included in the activity measurements for 'Other Radionuclides' until 1999.
- (4) Discharges of tritium are presented in Graph 2.
- (5) Annual discharge limits are given in Table 1.

**Graph 2 - Tritium Activity Discharged per year into the Hamoaze**



Note: (1) The discharge of effluent for 1987 was recorded from June 1987 to December 1987 (i.e. after DRDL's first Radioactive Substances Act authorisation was issued).

(2) Graph 2 shows the elevated discharges of tritium to the Hamoaze since 2002, which are associated primarily with the discharge of certain submarine effluents, arising from discrete operations during their Deep Maintenance Period. The graph illustrates the intermittent nature of discharges from DRD to the Hamoaze, which vary depending on the work packages being undertaken on nuclear powered submarines at DRD.

(3) Annual discharge limits are given in Table 1.

Figure 1 – Comparison of radiation doses

